

254 (SEQ ID NO:359), Figure 256 (SEQ ID NO:361), Figure 258 (SEQ ID NO:363), Figure 260 (SEQ ID NO:365), Figure 262 (SEQ ID NO:367), Figure 264 (SEQ ID NO:369), Figure 266 (SEQ ID NO:371), Figure 268 (SEQ ID NO:373), Figure 270 (SEQ ID NO:375), Figure 272 (SEQ ID NO:377), Figure 274 (SEQ ID NO:379), Figure 276 (SEQ ID NO:381), Figure 278 (SEQ ID NO:387), Figure 280 (SEQ ID NO:389), Figure 282 (SEQ ID NO:394), Figure 284 (SEQ ID NO:399), Figure 286 (SEQ ID NO:401), Figure 288 (SEQ ID NO:403), Figure 290 (SEQ ID NO:408), Figure 292 (SEQ ID NO:410), Figure 294 (SEQ ID NO:412), Figure 296 (SEQ ID NO:414), Figure 298 (SEQ ID NO:416), Figure 300 (SEQ ID NO:418), Figure 302 (SEQ ID NO:420), Figure 304 (SEQ ID NO:422), Figure 306 (SEQ ID NO:424), Figure 308 (SEQ ID NO:495), Figure 310 (SEQ ID NO:497), Figure 312 (SEQ ID NO:499), Figure 314 (SEQ ID NO:501), Figure 316 (SEQ ID NO:503), Figure 318 (SEQ ID NO:505), Figure 320 (SEQ ID NO:507), Figure 322 (SEQ ID NO:509), Figure 324 (SEQ ID NO:511), Figure 326 (SEQ ID NO:513), Figure 328 (SEQ ID NO:515) or Figure 330 (SEQ ID NO:517), lacking its associated signal peptide.

29. A method of detecting a PRO943 polypeptide in a sample suspected of containing a PRO943 polypeptide, said method comprising contacting said sample with a PRO183, PRO184 or PRO185 polypeptide and determining the formation of a PRO943/PRO183, PRO184 or PRO185 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO943 polypeptide in said sample.

30. The method according to Claim 29, wherein said sample comprises cells suspected of expressing said PRO943 polypeptide.

31. The method according to Claim 29, wherein said PRO183, PRO184 or PRO185 polypeptide is labeled with a detectable label.

32. The method according to Claim 29, wherein said PRO183, PRO184 or PRO185 polypeptide is attached to a solid support.

33. A method of detecting a PRO183, PRO184 or PRO185 polypeptide in a sample suspected of containing a PRO183, PRO184 or PRO185 polypeptide, said method comprising contacting said sample with a PRO943 polypeptide and determining the formation of a PRO943/PRO183, PRO184 or PRO185 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO183, PRO184 or PRO185 polypeptide in said sample.

34. The method according to Claim 33, wherein said sample comprises cells suspected of expressing said PRO183, PRO184 or PRO185 polypeptide.

35. The method according to Claim 33, wherein said PRO943 polypeptide is labeled with a detectable label.

36. The method according to Claim 33, wherein said PRO943 polypeptide is attached to a solid support.

37. A method of detecting a PRO331 polypeptide in a sample suspected of containing a PRO331 polypeptide, said method comprising contacting said sample with a PRO1133 polypeptide and determining the formation of a PRO331/PRO1133 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO331 polypeptide in said sample.

38. The method according to Claim 37, wherein said sample comprises cells suspected of expressing said PRO331 polypeptide.

39. The method according to Claim 37, wherein said PRO1133 polypeptide is labeled with a detectable label.

40. The method according to Claim 37, wherein said PRO1133 polypeptide is attached to a solid support.

41. A method of detecting a PRO1133 polypeptide in a sample suspected of containing a PRO1133 polypeptide, said method comprising contacting said sample with a PRO331 polypeptide and determining the formation of a PRO331/PRO1133 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO1133 polypeptide in said sample.

42. The method according to Claim 41, wherein said sample comprises cells suspected of expressing said PRO1133 polypeptide.

43. The method according to Claim 41, wherein said PRO331 polypeptide is labeled with a detectable label.

44. The method according to Claim 41, wherein said PRO331 polypeptide is attached to a solid support.

45. A method of detecting a PRO363 or PRO5723 polypeptide in a sample suspected of containing a PRO363 or PRO5723 polypeptide, said method comprising contacting said sample with a PRO1387 polypeptide and determining the formation of a PRO363 or PRO5723/PRO1387 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO363

or PRO5723 polypeptide in said sample.

46. The method according to Claim 45, wherein said sample comprises cells suspected of expressing said PRO363 or PRO5723 polypeptide.

5 47. The method according to Claim 45, wherein said PRO1387 polypeptide is labeled with a detectable label.

48. The method according to Claim 45, wherein said PRO1387 polypeptide is attached to a solid support.

10 49. A method of detecting a PRO1387 polypeptide in a sample suspected of containing a PRO1387 polypeptide, said method comprising contacting said sample with a PRO363 or PRO5723 polypeptide and determining the formation of a PRO363 or PRO5723/PRO1387 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO1387 polypeptide in said sample.

15 50. The method according to Claim 49, wherein said sample comprises cells suspected of expressing said PRO1387 polypeptide.

20 51. The method according to Claim 49, wherein said PRO363 or PRO5723 polypeptide is labeled with a detectable label.

52. The method according to Claim 49, wherein said PRO363 or PRO5723 polypeptide is attached to a solid support.

25 53. A method of detecting a PRO1114 polypeptide in a sample suspected of containing a PRO1114 polypeptide, said method comprising contacting said sample with a PRO3301 or PRO9940 polypeptide and determining the formation of a PRO1114/PRO3301 or PRO9940 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO1114 polypeptide in said sample.

30 54. The method according to Claim 53, wherein said sample comprises cells suspected of expressing said PRO1114 polypeptide.

35 55. The method according to Claim 53, wherein said PRO3301 or PRO9940 polypeptide is labeled with a detectable label.

56. The method according to Claim 53, wherein said PRO3301 or PRO9940 polypeptide is attached to a solid support.

57. A method of detecting a PRO3301 or PRO9940 polypeptide in a sample suspected of containing a PRO3301 or PRO9940 polypeptide, said method comprising contacting said sample with a PRO1114 polypeptide and determining the formation of a PRO3301 or PRO9940/PRO1114 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO3301 or PRO9940 polypeptide in said sample.

58. The method according to Claim 57, wherein said sample comprises cells suspected of expressing said PRO3301 or PRO9940 polypeptide.

59. The method according to Claim 57, wherein said PRO1114 polypeptide is labeled with a detectable label.

60. The method according to Claim 57, wherein said PRO1114 polypeptide is attached to a solid support.

61. A method of detecting a PRO1181 polypeptide in a sample suspected of containing a PRO1181 polypeptide, said method comprising contacting said sample with a PRO7170, PRO361 or PRO846 polypeptide and determining the formation of a PRO1181/PRO7170, PRO361 or PRO846 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO1181 polypeptide in said sample.

62. The method according to Claim 61, wherein said sample comprises cells suspected of expressing said PRO1181 polypeptide.

63. The method according to Claim 61, wherein said PRO7170, PRO361 or PRO846 polypeptide is labeled with a detectable label.

64. The method according to Claim 61, wherein said PRO7170, PRO361 or PRO846 polypeptide is attached to a solid support.

65. A method of detecting a PRO7170, PRO361 or PRO846 polypeptide in a sample suspected of containing a PRO7170, PRO361 or PRO846 polypeptide, said method comprising contacting said sample with a PRO1181 polypeptide and determining the formation of a PRO1181/PRO7170, PRO361 or PRO846 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO7170, PRO361 or PRO846 polypeptide in said sample.

66. The method according to Claim 65, wherein said sample comprises cells suspected of expressing said PRO7170, PRO361 or PRO846 polypeptide.

67. The method according to Claim 65, wherein said PRO1181 polypeptide is labeled with a detectable label.

68. The method according to Claim 65, wherein said PRO1181 polypeptide is attached to a solid support.

69. A method of linking a bioactive molecule to a cell expressing a PRO943 polypeptide, said method comprising contacting said cell with a PRO183, PRO184 or PRO185 polypeptide that is bound to said bioactive molecule and allowing said PRO943 and PRO183, PRO184 or PRO185 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

70. The method according to Claim 69, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

71. The method according to Claim 69, wherein said bioactive molecule causes the death of said cell.

72. A method of linking a bioactive molecule to a cell expressing a PRO183, PRO184 or PRO185 polypeptide, said method comprising contacting said cell with a PRO943 polypeptide that is bound to said bioactive molecule and allowing said PRO943 and PRO183, PRO184 or PRO185 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

73. The method according to Claim 72, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

74. The method according to Claim 73, wherein said bioactive molecule causes the death of said cell.

75. A method of linking a bioactive molecule to a cell expressing a PRO331 polypeptide, said method comprising contacting said cell with a PRO1133 polypeptide that is bound to said bioactive molecule and allowing said PRO331 and PRO1133 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

76. The method according to Claim 75, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

77. The method according to Claim 75, wherein said bioactive molecule causes the death of said cell.

78. A method of linking a bioactive molecule to a cell expressing a PRO1133 polypeptide, said method comprising contacting said cell with a PRO331 polypeptide that is bound to said bioactive molecule and allowing said PRO331 and PRO1133 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

79. The method according to Claim 78, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

80. The method according to Claim 78, wherein said bioactive molecule causes the death of said cell.

81. A method of linking a bioactive molecule to a cell expressing a PRO1387 polypeptide, said method comprising contacting said cell with a PRO363 or PRO5723 polypeptide that is bound to said bioactive molecule and allowing said PRO1387 and PRO363 or PRO5723 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

82. The method according to Claim 81, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

83. The method according to Claim 81, wherein said bioactive molecule causes the death of said cell.

84. A method of linking a bioactive molecule to a cell expressing a PRO363 or PRO5723 polypeptide, said method comprising contacting said cell with a PRO1387 polypeptide that is bound to said bioactive molecule and allowing said PRO1387 and PRO363 or PRO5723 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

85. The method according to Claim 84, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

86. The method according to Claim 84, wherein said bioactive molecule causes the death of said cell.

87. A method of linking a bioactive molecule to a cell expressing a PRO1114 polypeptide, said method comprising contacting said cell with a PRO3301 or PRO9940 polypeptide that is bound to said bioactive molecule and allowing said PRO1114 and PRO3301 or PRO9940 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

5 88. The method according to Claim 87, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

89. The method according to Claim 87, wherein said bioactive molecule causes the death of said cell.

10 90. A method of linking a bioactive molecule to a cell expressing a PRO3301 or PRO9940 polypeptide, said method comprising contacting said cell with a PRO1114 polypeptide that is bound to said bioactive molecule and allowing said PRO1114 and PRO3301 or PRO9940 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

15 91. The method according to Claim 90, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

20 92. The method according to Claim 90, wherein said bioactive molecule causes the death of said cell.

25 93. A method of linking a bioactive molecule to a cell expressing a PRO1181 polypeptide, said method comprising contacting said cell with a PRO7170, PRO361 or PRO846 polypeptide that is bound to said bioactive molecule and allowing said PRO1181 and PRO7170, PRO361 or PRO846 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

94. The method according to Claim 93, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

30 95. The method according to Claim 93, wherein said bioactive molecule causes the death of said cell.

35 96. A method of linking a bioactive molecule to a cell expressing a PRO7170, PRO361 or PRO846 polypeptide, said method comprising contacting said cell with a PRO1181 polypeptide that is bound to said bioactive molecule and allowing said PRO1181 and PRO7170, PRO361 or PRO846 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

97. The method according to Claim 96, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

98. The method according to Claim 96, wherein said bioactive molecule causes the death of said cell.

99. A method of modulating at least one biological activity of a cell expressing a PRO943 polypeptide, said method comprising contacting said cell with a PRO183, PRO184 or PRO185 polypeptide or an anti-PRO943 antibody, whereby said PRO183, PRO184 or PRO185 polypeptide or said anti-PRO943 antibody binds to said PRO943 polypeptide, thereby modulating at least one biological activity of said cell.

100. The method according to Claim 99, wherein said cell is killed.

101. A method of modulating at least one biological activity of a cell expressing a PRO183, PRO184 or PRO185 polypeptide, said method comprising contacting said cell with a PRO943 polypeptide or an anti-PRO183, anti-PRO184 or anti-PRO185 antibody, whereby said PRO943 polypeptide or said anti-PRO183, anti-PRO184 or anti-PRO185 antibody binds to said PRO183, PRO184 or PRO185 polypeptide, thereby modulating at least one biological activity of said cell.

102. The method according to Claim 101, wherein said cell is killed.

103. A method of modulating at least one biological activity of a cell expressing a PRO331 polypeptide, said method comprising contacting said cell with a PRO1133 polypeptide or an anti-PRO331 antibody, whereby said PRO1133 polypeptide or said anti-PRO331 antibody binds to said PRO331 polypeptide, thereby modulating at least one biological activity of said cell.

104. The method according to Claim 103, wherein said cell is killed.

105. A method of modulating at least one biological activity of a cell expressing a PRO1133 polypeptide, said method comprising contacting said cell with a PRO331 polypeptide or an anti-PRO1133 antibody, whereby said PRO331 polypeptide or said anti-PRO1133 antibody binds to said PRO1133 polypeptide, thereby modulating at least one biological activity of said cell.

106. The method according to Claim 105, wherein said cell is killed.



107. A method of modulating at least one biological activity of a cell expressing a PRO1387 polypeptide, said method comprising contacting said cell with a PRO363 or PRO5723 polypeptide or an anti-PRO1387 antibody, whereby said PRO363 or PRO5723 polypeptide or said anti-PRO1387 antibody binds to said PRO1387 polypeptide, thereby modulating at least one biological activity of said cell.

5 108. The method according to Claim 107, wherein said cell is killed.

109. A method of modulating at least one biological activity of a cell expressing a PRO363 or PRO5723 polypeptide, said method comprising contacting said cell with a PRO1387 polypeptide or an anti-PRO363 or anti-PRO5723 antibody, whereby said PRO1387 polypeptide or said anti-PRO363 or anti-PRO5723 antibody binds to said PRO363 or PRO5723 polypeptide, thereby modulating at least one biological activity of said cell.

110. The method according to Claim 109, wherein said cell is killed.

15 111. A method of modulating at least one biological activity of a cell expressing a PRO1114 polypeptide, said method comprising contacting said cell with a PRO3301 or PRO9940 polypeptide or an anti-PRO1114 antibody, whereby said PRO3301 or PRO9940 polypeptide or said anti-PRO1114 antibody binds to said PRO1114 polypeptide, thereby modulating at least one biological activity of said cell.

20 112. The method according to Claim 111, wherein said cell is killed.

25 113. A method of modulating at least one biological activity of a cell expressing a PRO3301 or PRO9940 polypeptide, said method comprising contacting said cell with a PRO1114 polypeptide or an anti-PRO3301 or anti-PRO9940 antibody, whereby said PRO1114 polypeptide or said anti-PRO3301 or anti-PRO9940 antibody binds to said PRO3301 or PRO9940 polypeptide, thereby modulating at least one biological activity of said cell.

114. The method according to Claim 113, wherein said cell is killed.

30 115. A method of modulating at least one biological activity of a cell expressing a PRO1181 polypeptide, said method comprising contacting said cell with a PRO7170, PRO361 or PRO846 polypeptide or an anti-PRO1181 antibody, whereby said PRO7170, PRO361 or PRO846 polypeptide or said anti-PRO1181 antibody binds to said PRO1181 polypeptide, thereby modulating at least one biological activity of said cell.

35 116. The method according to Claim 115, wherein said cell is killed.

117. A method of modulating at least one biological activity of a cell expressing a PRO7170, PRO361 or PRO846 polypeptide, said method comprising contacting said cell with a PRO1181 polypeptide or an anti-PRO7170, anti-PRO361 or anti-PRO846 antibody, whereby said PRO1181 polypeptide or said anti-PRO7170, anti-PRO361 or anti-PRO846 antibody binds to said PRO7170, PRO361 or PRO846 polypeptide, thereby modulating at least one biological activity of said cell.

5

118. The method according to Claim 117, wherein said cell is killed.